

## Module C Comprehensive

- Verify the on-line record [the grades page]
- No review period
  - But encouraged to discuss it with the instructor **No negotiation**
- Return the package today
  - May take home if returned by coming Tue.
  - Available for review in the instructor's office
- Mini Project Phase 2
  - Review and include it in the Final Package

CMSC210 D2

1

## Unit D2: Relations

### Today

- Understand and use **equivalence relation**, **equivalence class**, and **partition**
- Prove certain properties of relations, etc.
- Take-home exercises
  - Space stations, Bogus proofs & paradoxes

CMSC210 D2

2

### Section 1

## Example Relations


- Weeding
  - Relation: "is connected to (through the root system)"
- Academic integrity
  - Relation: "shares information with"
- Room temperature (temp. zones)
  - Relation: "has opening to"
- Flash Psychic **Properties?**

CMSC210 D2

3

## Equivalence Relation

- Relation with all of the following properties:
  - Reflexive
  - Symmetric
  - Transitive
- Cf. the **relation** in a poset




What can you say about the set members?

CMSC210 D2

4

## Equivalence Class

- A subset that are mutually related through an equivalence relation
  - Notation:  $[x]$  (equivalence class including  $x$ )



$$[a] = [b] = [c] = \{a, b, c\}$$

$$[d] = [e] = \{d, e\}$$


- (more formally) For an **equivalence relation**  $R \subseteq A \times A$ ,  $X$  is an **equivalence class** if  $X \subseteq A$  and  $\forall x, y \in X ((x, y) \in R)$

CMSC210 D2

5

## Partition

- The set of equivalence classes




$$\{\{a, b, c\}, \{d, e\}\} = \{[a], [d]\}$$

- Properties
  - The subsets are all disjoint.
  - The union of the subsets covers the entire set.
- (more formally) For an **equivalence relation**  $R \subseteq A \times A$ , the **partition** of  $A$  is defined as  $\{[x] \mid x \in A\}$

CMSC210 D2

6

## Group Exercise 1: Weeding



- Define the equivalence relation
- Find all equivalence classes
- Find the partition

CMSC210 D2

7

## Group Exercise 2: Flash Psychic

- Informally describe the involved equivalent relation  $R$  between the numbers on the number-symbol correspondence charts
- Formally define the partition induced by  $R$

CMSC210 D2

8

## Section Summary

- Equivalence relation
  - Reflexive, symmetric, transitive relation
- Equivalence relation forms equivalence classes.
- Equivalence relation **induces** a partition

CMSC210 D2

9

## Section 2

## Mathemagic

- Question: Is 0.999... equal to 1.0?

Proof?

CMSC210 D2

10

## Proof Structure

Review: B5

Hypothesis:  $o \text{ ® } f, o$

Proof of  $f$  from the above hypothesis

1.  $o \text{ ® } f$  [hyp]
  2.  $o$  [hyp]
  3.  $f$  [MP: 1, 2]
- conclusion/theorem
- justification for each step (must be semantically correct)
- line numbering

CMSC210 D2

11

## Generalized Flash Psychic

- Hypotheses
  - $m$  is a positive integer.
  - $R = \{(x, y) \mid (x \bmod m) = (y \bmod m)\}$
- Conclusion:  $R$  is an equivalence relation. Example members?
- Note:  $x \bmod y$  is the remainder of  $x$  divided by  $y$ . Cf. FSA to recognize 10, 010, 110, 0010, 0110, 1010, 1110

CMSC210 D2

12

## Proof

1.  $R$  is reflexive. [Lemma 1]
2.  $R$  is symmetric. [Lemma 2]
3.  $R$  is transitive. [Lemma 3]
4.  $R$  is an equivalence relation. [Def. eq. rel. 1.-3.]

Note: **Lemma** = preliminary theorem

## Lemma 1: $R$ is reflexive.

Hypotheses

- $m$  is a positive integer.
- $R = \{(x, y) \mid (x \bmod m) = (y \bmod m)\}$

Conclusion: For any  $x$ ,  $(x, x) \in R$

1.  $(x \bmod m) = (x \bmod m)$  [Def. function]
2.  $R$  is reflexive. [Def. reflexive]

"trivial," "obvious," "easy," etc.?

## Lemma 2: $R$ is symmetric.

Def: For any  $x, y$ , if  $(x, y) \in R$ , then  $(y, x) \in R$

Hypotheses

- $m$  is a positive integer.
- $R = \{(x, y) \mid (x \bmod m) = (y \bmod m)\}$
- $(x, y) \in R$

Conclusion:  $(y, x) \in R$

1.  $(x \bmod m) = (y \bmod m)$  [Hyp.]
2.  $(y \bmod m) = (x \bmod m)$  [\_\_\_\_\_]
3.  $R$  is symmetric. [Def. symmetric]

## Lemma 3: $R$ is transitive.

Def: For any  $x, y, z$ , if  $(x, y), (y, z) \in R$ , then  $(x, z) \in R$

Hypotheses

- $m$  is a positive integer.
- $R = \{(x, y) \mid (x \bmod m) = (y \bmod m)\}$
- $(x, y), (y, z) \in R$

Conclusion:  $(x, z) \in R$

1.  $(x \bmod m) = (y \bmod m)$  [Hyp.]
2.  $(y \bmod m) = (z \bmod m)$  [Hyp.]
3. \_\_\_\_\_ [\_\_\_\_\_]
4.  $R$  is transitive. [Def. transitive]

## Prove

- There is a country where the breakfast is served with boiled eggs cooked for either 5 or 8 minutes.

## Disprove

- There is a unicorn in this room.

## Prove or Disprove


- There is a cave system whose total length reaches hundreds of miles.
  - Imagine a tunnel from Philadelphia to, say, Boston

CMSC210 D2

19

## Prove or Disprove

- Nondeterminism can be simulated as determinism applying the idea of “power set.”
- Note: Try this in terms of FSA's



CMSC210 D2

20

## Proof/Disproof Strategy

- To **prove** a **general** statement, follow the proof pattern discussed in Unit B5. That is, identify the hypotheses (if any) and the conclusion, number proof steps, and give justification for each step.
- To **disprove** a **general** statement, give a counterexample.
- To **prove** an **existential** statement, give an example [proof by existence/example].
- To **disprove** an **existential** statement, follow the proof pattern as noted above and justify that there is **no** such thing.
- If not sure to prove/disprove, try both.

CMSC210 D2

21

Review: B5

## Proof Methods Summary

- **Direct proof:** Straightforward proof of a conclusion from hypotheses (if any)
  - **Proof by cases:** Prove exclusive and exhaustive cases separately
- **Indirect proof:** Uses a proof of a proposition different from the conclusion
  - **Proof by contradiction:** Assume the negation of the conclusion and derive a contradiction

CMSC210 D2

22

## Section Summary

- Roles of logic
  - To specify structures
  - To justify by proving or disproving (semantically correct syntactic manipulation)
- Approaches to proof
  - General statement (formally,  $\forall x \dots$ )
    - Prove: Cover all possibilities
    - Disprove: Counterexample
  - Existential statement (formally,  $\exists x \dots$ )
    - Prove: Example
    - Disprove: Cover all impossibilities

CMSC210 D2

23

## Summary Exercise

- **Hypothesis:** Direct or indirect interaction between objects in the universe is an equivalence relation. *Make up your own universe.*
- **Prove or disprove:** The universe is a partition (induced by the above equivalence relation) with a cardinality greater than 1.
  - If difficult, explain why (definition? proof step?).
- **[optional]** Philosophical implications of your answer?
- **Questions/Comments/Suggestions**

CMSC210 D2

24